

Prescription Label

Patient Name:

Species:

Drug Name & Strength:

Directions (amount to give how often & for how long):

Prescribing Veterinarian's Name & Contact Information:

Refills:

[Content to be provided by prescribing veterinarian]

Famciclovir

(fam-sye-klow-veer)

Description: Antiviral (Herpes)

Other Names for this Medication: Famvir®

Common Dosage Forms: Veterinary: None. **Human:** 125 mg, 250 mg, & 500 mg tablets.

This information sheet does not contain all available information for this medication. It is to help answer commonly asked questions and help you give the medication safely and effectively to your animal. If you have other questions or need more information about this medication, contact your veterinarian or pharmacist.

Key Information

- ▶ Medicine used to control, but not cure, herpesvirus infections in cats.
- ▶ Try giving medicine without food, but if your cat vomits or acts sick after getting the medicine, give with a small amount of food. Because there is limited experience with this drug in veterinary medicine, be sure to report any concerns about your cat to your veterinarian while your cat is taking this medicine.
- ▶ Usually well tolerated.

How is this medication useful?

Famciclovir is an antiviral agent used to control, but not cure, herpesvirus infections in cats. The FDA (U.S. Food & Drug Administration) has approved famciclovir for use in humans, but it is not officially approved for use in animals. The FDA allows veterinarians to prescribe and use human products containing famciclovir in animals in certain situations. You and your veterinarian can discuss why this drug is the most appropriate choice.

What should I tell my veterinarian to see if this medication can be safely given?

Many things might affect how well famciclovir will work in your animal. Be sure to discuss the following with your veterinarian so together you can make the best treatment decisions.

- ▶ Other drugs can interact with famciclovir, so be sure to tell your veterinarian and pharmacist what medications (including vitamins, supplements, or herbal therapies) you give your animal, including the amount and time you give each.
- ▶ Tell your veterinarian about any conditions or diseases your animal may have now or has had in the past.
- ▶ If your animal has been treated for the same disease or condition in the past, tell your veterinarian about the treatment and how well it did or did not work.

- ▶ If your animal is pregnant or nursing, talk to your veterinarian about the risks of using this drug.
- ▶ Tell your veterinarian and pharmacist about any medication side effects (including allergic reactions, lack of appetite, diarrhea, itching, hair loss) your animal has developed in the past.

How long until I will know if this medication is working, and how long will the effects of this medication last?

This medication takes several doses before you will see any benefits. The medication's full effects may not be evident until it has been given consistently for several weeks, although you may see some side effects (eg, upset stomach or sleepiness) immediately. Your animal's clinical signs will improve gradually over the time the medication takes to become fully effective.

The effects of this medication are short-lived, meaning they will stop working within 24 hours, although the benefits may be prolonged if your animal has decreased kidney and/or liver function.

When should this medication not be used or be used very carefully?

No drug is 100% safe in all patients, but your veterinarian will discuss with you any specific concerns about using famciclovir in your animal.

This drug **SHOULD NOT** be used in patients:

- ▶ That are allergic to it.

This drug should be used **WITH CAUTION** in patients:

- ▶ That have kidney problems.

If your animal has any of these conditions, talk to your veterinarian about the potential risks versus benefits.

What are the side effects of this medication?

Side effects that usually are not serious include:

- ▶ Decreased appetite.
- ▶ Increased amounts of urine.

If any of these signs are severe, worsen, or continue to be a problem, contact your veterinarian.

If my animal gets too much of this medication (an overdose), what should I do?

If you witness or suspect an overdose, contact your veterinarian or an animal poison control center for further advice. Animal poison control centers open 24 hours a day include: **Pet Poison HELPLINE** (855-764-7661) and **ASPCA Animal Poison Control Center** (888-426-4435); a consultation fee may be charged for these services.

How should this medication be given?

For famciclovir to work properly, give it exactly as your veterinarian has prescribed. Check the prescription label to be sure you are giving the drug correctly.

- ▶ The drug may be given either with food or on an empty stomach. If your animal vomits or acts sick after receiving the drug on an empty stomach, try giving the next dose with food or a small treat. If vomiting continues, contact your veterinarian.
- ▶ Compounded liquid forms of famciclovir must be measured carefully. Your veterinarian or pharmacist can help by providing special measuring spoons or syringes.
- ▶ If you have difficulty getting your animal to take the medicine, contact your veterinarian or pharmacist for tips to help with dosing and reduce the stress of medication time for both you and your animal.
- ▶ Famciclovir can be given for various lengths of time. Be sure you understand how long your veterinarian wants you to continue giving famciclovir. Prescription refills may be necessary before the therapy will be complete. If you want to stop giving this medication, talk to your veterinarian, as there may be important reasons to continue giving it.

What should I do if I miss giving a dose of this medication?

If you miss a dose, give it when you remember, but if it is close enough to the time for the next dose, skip the dose you missed and give it at the next scheduled time. After that, return to the regular dosing schedule. Do not double-up or give extra doses.

How should I store this medication?

- ▶ Store this medication in the original prescription bottle or an approved dosage reminder container (ie, pill minder) at room temperature and protected from light.
- ▶ If your veterinarian or pharmacist has prepared (compounded) a special formulation for your animal, follow the storage recommendations and expiration date for the product.
- ▶ Keep away from children and other animals.

Can handling this medication be hazardous to me, my family, or other animals?

There are no specific precautions required when handling this medication unless you are allergic to it. Wash your hands after handling any medication.

How should I dispose of this medication if I don't use it all?

- ▶ Do not flush this medication down the toilet or wash it down the sink. If a community drug “take-back” program is available, use this option. If there is no take-back program, mix the drug with coffee grounds or cat litter (to make it undesirable to children and animals and unrecognizable to people who might go through your trash), place the mixture in a sealable plastic bag to keep it from leaking out, and throw the bag out with the regular trash.
- ▶ Do not save leftover medication for future use or give it to others to use.

What other information is important for this medication?

- ▶ Use of famciclovir may not be allowed in certain animal competitions. Check rules and regulations before entering your animal in a competition while this medication is being administered.
- ▶ If your animal needs to be on this medicine for long periods of time, your veterinarian may recommend periodic blood tests to ensure your animal's body is tolerating the medicine well. Do not miss these important follow-up visits.

If you have any other questions about this medication, contact your veterinarian or pharmacist.

Special Instructions:
